

Product References

Poietics® Preadipocyte Cells

Cells

1. *Aoki S, Toda S, Ando T, Sugihara H. Bone marrow stromal cells, preadipocytes, and dermal fibroblasts promote epidermal regeneration in their distinctive fashions. *Mol Biol Cell*. 2004 Oct; 15(10): 4647-57.
2. Benson SC, Pershadsingh HA, Ho CI, Chittiboyina A, Desai P, Pravenec M, Qi N, Wang J, Avery MA, Kurtz TW. Identification of telmisartan as a unique angiotensin II receptor antagonist with selective PPAR γ – modulating activity. *Hypertension*. 2004; 43: 993.
3. Fernández-Veledo S, Vila-Bedmar R, Nieto-Vazquez I, Lorenzo M. J Clin Endocrinol Metab. c-Jun N-terminal kinase 1/2 activation by tumor necrosis factor- α induces insulin resistance in human visceral but not subcutaneous adipocytes: reversal by liver X receptor agonists. *J Clin Endocrinol Metab*. 2009 Sep;94(9):3583-93. Epub 2009 Jun 30.
4. Gazi E, Dwyer J, Lockyer NP, Gardner P, Shanks JH, Roulson J, Hart CA, Clarke NW, Brown MD. Biomolecular profiling of metastatic prostate cancer cells in bone marrow tissue using FTIR microspectroscopy: a pilot study. *Anal Bioanal Chem*. 2007 Mar;387(5):1621-31.
5. *Grohmann M, Sabin M, Holly J, Shield J, Crowne E, Stewart C. Characterization of differentiated subcutaneous and visceral adipose tissue from children: the influences of TNF- α and IGF-I. *J Lipid Res*. 2005 Jan; 46(1): 93-103.
6. Héту PO, Riendeau D. Down-regulation of microsomal prostaglandin E2 synthase-1 in adipose tissue by high-fat feeding. *Obesity (Silver Spring)*. 2007 Jan;15(1):60-8.
7. Hirano T, Morii H, Nakazawa K, Murakami D, Yamashita A, Asahi J, Orimo H, Tamae K, Tokura Y. Enhancement of adipogenesis induction by conditioned media obtained from cancer cells. *Cancer Letters*. 2008. Ahead of Print.
8. Hong KM, Burdick MD, Phillips RJ, Heber D, Strieter RM. Characterization of human fibrocytes as circulating adipocyte progenitors and the formation of human adipose tissue in SCID mice. *FASEB J*. 2005 Dec; 19(14):2029-31.
9. *Hutley LJ, Newell FM, Joyner JM, Suchting SJ, Herington AC, Cameron DP, Prins JB. Effects of rosiglitazone and linoleic acid on human preadipocyte differentiation. *Eur J Clin Invest*. 2003 Jul; 33(7): 574-81.
10. Kobashi C, Asamizu S, Ishiki M, Iwata M, Usui I, Yamazaki K, Tobe K, Kobayashi M, Urakaze M. Inhibitory effect of IL-8 on insulin action in human adipocytes via MAP kinase pathway. *J Inflamm (Lond)*. 2009 Aug 27;6:25.
11. Kohlstedt K, Gershon C, Trouvain C, Hofmann WK, Fichtlscherer S, Fleming I. Angiotensin-converting enzyme (ACE) inhibitors modulate cellular retinol-binding protein 1 and adiponectin expression in adipocytes via the ACE-dependent signaling cascade. *Mol Pharmacol*. 2009 Mar;75(3):685-92. Epub 2008 Dec 29.
12. Onuma M, Bub JD, Rummel TL, Iwamoto Y. Prostate cancer cell-adipocyte interaction. *J Biol Chem*. 2003; 278(43): 42660-7.
13. Qi L, Kang K, Zhang C, van Dam RM, Kraft P, Hunter D, Lee CH, Hu FB. Fat mass-and obesity-associated (FTO) gene variant is associated with obesity: longitudinal analyses in two cohort studies and functional test. *Diabetes*. 2008 Nov;57(11):3145-51.
14. Ren N, Kaplan R, Hernandez M, Cheng K, Jin L, Taggart AK, Zhu AY, Gan X, Wright SD, Cai TQ. Phenolic acids suppress

- adipocyte lipolysis via activation of the nicotinic acid receptor GPR109A (HM74a/PUMA-G). *J Lipid Res.* 2009 May;50(5):908-14. Epub 2009 Jan 9.
15. *Tchkonia T, Tchoukalova YD, Giorgadze N, Pirtskhalava T, Karagiannides I, Forse RA, Koo A, Stevenson M, Chinnappan D, Cartwright A, Jensen MD, Kirkland JL. Abundance of two human preadipocyte subtypes with distinct capacities for replication, adipogenesis, and apoptosis varies among fat depots. *Am J Physiol Endocrinol Metab.* 2005; 288(1): E267-77.
 16. Wang M, Crisostomo PR, Herring C, Meldrum KK, Meldrum DR. Human progenitor cells from bone marrow or adipose tissue produce VEGF, HGF, and IGF-I in response to TNF by a p38 MAPK-dependent mechanism. *Am J Physiol Regul Integr Comp Physiol.* 2006; 291(4):R880-4.
 17. Wang M, Tsai BM, Crisostomo PR, Meldrum DR. Pretreatment with adult progenitor cells improves recovery and decreases native myocardial proinflammatory signaling after ischemia. *Shock.* 2006 May;25(5):454-9.
 18. Wolk K, Haugen HS, Xu W, Witte E, Waggle K, Anderson M, Vom Baur E, Witte K, Warszawska K, Philipp S, Johnson-Leger C, Volk HD, Sterry W, Sabat R. IL-22 and IL-20 are key mediators of the epidermal alterations in psoriasis while IL-17 and IFN-gamma are not. *J Mol Med.* 2009 May;87(5):523-36. Epub 2009 Mar 30.
 19. Yamagishi SI, Matsui T, Adachi H, Takeuchi M. Positive association of circulating levels of advanced glycation end products (AGEs) with pigment epithelium-derived factor (PEDF) in a general population. *Pharmacol Res.* 2009 Jul 25.
 20. Yamazaki Y, Kawano Y, Uebayasi M. Induction of adiponectin by natural and synthetic phenolamides in mouse and human preadipocytes and its enhancement by docosahexaenoic acid. *Life Sci.* 2008 Jan 30;82(5-6):290-300.
 21. Yang M, Sun J, Zhang T, Liu J, Zhang J, Shi MA, Darakhshan F, Guerre-Millo M, Clement K, Gelb BD, Dolgiov G, Shi GP. Deficiency and inhibition of cathepsin K reduce body weight gain and increase glucose metabolism in mice. *Arterioscler Thromb Vasc Biol.* 2008 Dec;28(12):2202-8.
 1. Ambati S, Kim HK, Yang JY, Lin J, Della-Fera MA, Baile CA. Effects of leptin on apoptosis and adipogenesis in 3T3-L1 adipocytes. *Biochem Pharmacol.* 2007 Feb 1;73(3):378-84.
 2. Benhaddou-Andaloussi A, Martineau LC, Spoor D, Vuong T, Leduc C, Joly E, Burt A, Meddah B, Settaf A, Arnason JT, Prentki M, Haddad PS. Antidiabetic Activity of *Nigella sativa* Seed Extract in Cultured Pancreatic β -cells, Skeletal Muscle Cells, and Adipocytes. *Pharmaceutical Biology.* 2008 Jan; 46(1 & 2): 96-104.
 3. Carfi' M, Croera C, Ferrario D, Campi V, Bowe G, Pieters R, Gribaldo L. TBTC induces adipocyte differentiation in human bone marrow long term culture. *Toxicology.* 2008 Jul 10;249(1):11-8.
 4. Einstein M, Akiyama TE, Castriota GA, Wang CF, McKeever B, Mosley RT, Becker JW, Moller DE, Meinke PT, Wood HB, Berger JP. The differential interactions of peroxisome proliferator-activated receptor gamma ligands with Tyr473 is a physical basis for their unique biological activities. *Mol Pharmacol.* 2008 Jan;73(1):62-74.
 5. Forte G, Carotenuto F, Pagliari F, Pagliari S, Cossa P, Fiaccavento R, Ahluwalia A, Vozzi G, Vinci B, Serafino A, Rinaldi A, Traversa E, Carosella L, Minieri M, Di Nardo P. Criticality of the biological and physical stimuli array inducing resident cardiac stem cell determination. *Stem Cells.* 2008 Aug;26(8):2093-103.
 6. Fujita T, Furukawa S, Morita K, Ishihara T, Shiotani M, Matsushita Y, Matsuda M, Shimomura I. Glucosamine induces lipid accumulation and adipogenic change in C2C12 myoblasts. *Biochem Biophys Res Commun.* 2005 Mar 11;328(2):369-74.
 7. Gonda K, Shigeura T, Sato T, Matsumoto D, Suga H, Inoue K, Aoi N, Kato H, Sato K, Murase S, Koshima I, Yoshimura K. Preserved proliferative capacity and multipotency of human adipose-derived stem cells after long-term cryopreservation. *Plast Reconstr Surg.* 2008 Feb; 121(2):401-10.
 8. Gupta V, Bhasin S, Guo W, Singh R, Miki R, Chauhan P, Choong K, Tchkonia T, Lebrasseur NK, Flanagan JN, Hamilton JA, Viereck JC, Narula NS, Kirkland JL, Jasuja

- R. Effects of dihydrotestosterone on differentiation and proliferation of human mesenchymal stem cells and preadipocytes. *Mol Cell Endocrinol*. 2008 Dec 16;296(1-2):32-40.
9. Hausman DB, Park HJ, Hausman GJ. Isolation and culture of preadipocytes from rodent white adipose tissue. *Methods Mol Biol*. 2008;456:201-19.
 10. Hirano T, Morii H, Nakazawa K, Murakami D, Yamashita A, Asahi J, Orimo H, Tamae K, Tokura Y. Enhancement of adipogenesis induction by conditioned media obtained from cancer cells. *Cancer Letters*. 2008. Ahead of Print.
 11. Liang J, Fu M, Ciociola E, Chandalia M, Abate N. Role of ENPP1 on adipocyte maturation. *PLoS ONE*. 2007 Sep 12;2(9):e882.
 12. Marshak DR, Greenwalt DE. Differentiating primary human cells in rapid-throughput discovery applications. *Methods Mol Biol*. 2007;356:121-8.
 13. Martineau LC, Couture A, Spoor D, Benhaddou-Andaloussi A, Harris C, Meddah B, Leduc C, Burt A, Vuong T, Mai Le P, Prentki M, Bennett SA, Arnason JT, Haddad PS. Anti-diabetic properties of the Canadian lowbush blueberry *Vaccinium angustifolium* Ait. *Phytomedicine*. 2006 Nov;13(9-10):612-23.
 14. Matsubara Y, Saito E, Suzuki H, Watanabe N, Murata M, Ikeda Y. Generation of megakaryocytes and platelets from human subcutaneous adipose tissues. *Biochem Biophys Res Commun*. 2009 Jan 23;378(4):716-20.
 15. Matsumoto T, Kano K, Kondo D, Fukuda N, Iribe Y, Tanaka N, Matsubara Y, Sakuma T, Satomi A, Otaki M, Ryu J, Mugishima H. Mature adipocyte-derived dedifferentiated fat cells exhibit multilineage potential. *J Cell Physiol*. 2008 Apr;215(1):210-22.
 16. Meng X, Ichim TE, Zhong J, Rogers A, Yin Z, Jackson J, Wang H, Ge W, Bogin V, Chan KW, Thébaud B, Riordan NH. Endometrial regenerative cells: a novel stem cell population. *J Transl Med*. 2007 Nov 15;5:57.
 17. Mitchell F, Scanlan T, Taylor P. A role for 3-iodothyronamine in regulation of adipocyte lipolysis. *Endocrine Abstracts*. 2008; 15: 380.
 18. Muthusamy VS, Anand S, Sangeetha KN, Sujatha S, Arun B, Lakshmi BS. Tannins present in *Cichorium intybus* enhance glucose uptake and inhibit adipogenesis in 3T3-L1 adipocytes through PTP1B inhibition. *Chem Biol Interact*. 2008 Jul 10;174(1):69-78.
 19. Park HJ, Yang JY, Ambati S, Della-Fera MA, Hausman DB, Rayalam S, Baile CA. Combined effects of genistein, quercetin, and resveratrol in human and 3T3-L1 adipocytes. *J Med Food*. 2008 Dec;11(4):773-83.
 20. Rayalam S, Della-Fera MA, Ambati S, Boyan B, Baile CA. Enhanced effects of guggulsterone plus 1,25(OH)2D3 on 3T3-L1 adipocytes. *Biochem Biophys Res Commun*. 2007 Dec 21;364(3):450-6.
 21. Rayalam S, Della-Fera MA, Yang JY, Park HJ, Ambati S, Baile CA. Resveratrol potentiates genistein's antiadipogenic and proapoptotic effects in 3T3-L1 adipocytes. *J Nutr*. 2007 Dec; 137(12):2668-73.
 22. Ricchi M, Odoardi MR, Carulli L, Anzivino C, Ballestri S, Pinetti A, Fantoni LI, Marra F, Bertolotti M, Banni S, Lonardo A, Carulli N, Loria P. Differential effect of oleic and palmitic acid on lipid accumulation and apoptosis in cultured hepatocytes. *J Gastroenterol Hepatol*. 2009 Jan 13. [Epub ahead of print]
 23. Rival Y, Stennevin A, Puech L, Rouquette A, cathala C, Lestienne F, Dupont-Passelaigue E, Patoiseau JF, Wurch T, Junquero D. Human Adipocyte fatty acid-binding protein (aP2) gene promoter-driven reporter assay discriminates nonlipogenic peroxisome proliferators-activated receptor gamma ligands. *J Pharmacol Exp Ther*. 2004 Nov; 311(2): 467-75.
 24. Shirakawa K, Maeda S, Gotoh T, Hayashi M, Shinomiya K, Ehata S, Nishimura R, Mori M, Onozaki K, Hayashi H, Uematsu S, Akira S, Ogata E, Miyazono K, Imamura T. CCAAT/enhancer-binding protein homologous protein (CHOP) regulates osteoblast differentiation. *Mol Cell Biol*. 2006 Aug;26(16):6105-16.
 25. Spoor DC, Martineau LC, Leduc C, Benhaddou-Andaloussi A, Meddah B, Harris C, Burt A, Fraser MH, Coonishish J, Joly E, Cuerrier A, Bennett SA, Johns T, Prentki M, Arnason JT, Haddad PS. Selected plant species from the Cree pharmacopoeia of

- northern Quebec possess anti-diabetic potential. *Can J Physiol Pharmacol*. 2006 Aug-Sep;84(8-9):847-58.
26. Suga H, Matsumoto D, Inoue K, Shigeura T, Eto H, Aoi N, Kato H, Abe H, Yoshimura K. Numerical measurement of viable and nonviable adipocytes and other cellular components in aspirated fat tissue. *Plast Reconstr Surg*. 2008 Jul;122(1):103-14.
 27. Suga H, Shigeura T, Matsumoto D, Inoue K, Kato H, Aoi N, Murase S, Sato K, Gonda K, Koshima I, Yoshimura K. Rapid expansion of human adipose-derived stromal cells preserving multipotency. *Cytotherapy*. 2007;9(8):738-45.
 28. Takenaga M, Ohta Y, Tokura Y, Hamaguchi Y, Igarashi R, Disrathakit A, Doi N. Lipid Microsphere Formulation Containing Rifampicin Targets Alveolar Macrophages. *Drug Delivery*. 2008 Mar; 15(3): 169 – 175.
 29. Tartaglia LA. Complementary new approaches enable repositioning of failed drug candidates. *Expert Opin Investig Drugs*. 2006 Nov;15(11):1295-8.
 30. Vuong T, Martineau LC, Ramassamy C, Matar C, Haddad PS. Fermented Canadian lowbush blueberry juice stimulates glucose uptake and AMP-activated protein kinase in insulin-sensitive cultured muscle cells and adipocytes. *Can J Physiol Pharmacol*. 2007 Sep;85(9):956-65.
 31. Yamakawa J, Ishigaki Y, Takano F, Takahashi T, Yoshida J, Moriya J, Takata T, Tatsuno T, Sasaki K, Ohta T, Takegami T, Yoshizaki F. The Kampo medicines Orengedokuto, Bofutsushosan and Boiogito have different activities to regulate gene expressions in differentiated rat white adipocytes: comprehensive analysis of genetic profiles. *Biol Pharm Bull*. 2008 Nov;31(11):2083-9.
 32. Yang JY, Della-Fera MA, Baile CA. Guggulsterone inhibits adipocyte differentiation and induces apoptosis in 3T3-L1 cells. *Obesity (Silver Spring)*. 2008 Jan;16(1):16-22.
 33. Yang JY, Della-Fera MA, Hartzell DL, Nelson-Dooley C, Hausman DB, Baile CA. Esculetin induces apoptosis and inhibits adipogenesis in 3T3-L1 cells. *Obesity (Silver Spring)*. 2006 Oct;14(10):1691-9
 34. Yang JY, Della-Fera MA, Rayalam S, Ambati S, Hartzell DL, Park HJ, Baile CA. Enhanced inhibition of adipogenesis and induction of apoptosis in 3T3-L1 adipocytes with combinations of resveratrol and quercetin. *Life Sciences*. 2008: Ahead of Print.
 35. Yang JY, Della-Fera MA, Rayalam S, Baile CA. Effect of xanthohumol and isoxanthohumol on 3T3-L1 cell apoptosis and adipogenesis. *Apoptosis*. 2007 Nov;12(11):1953-63.
 36. Yin Y, Yuan H, Wang C, Pattabiraman N, Rao M, Pestell RG, Glazer RI. 3-phosphoinositide-dependent protein kinase-1 activates the peroxisome proliferator-activated receptor-gamma and promotes adipocyte differentiation. *Mol Endocrinol*. 2006 Feb;20(2):268-78.

* References not specifically citing the use of Lonza cells, media, or reagents in their research.

+ Denotes sections containing only the articles published within the last ten years.